

AD-A114 697

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

L INFINITY-UPPER BOUND OF L2-PROJECTIONS ONTO SPLINES AT A GEOM--ETC(U)

FEB 82 R JIA

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2335

NL




END

DATE

10 MAR 82

6 42

DTIC



DAI 14697

2

MRC Technical Summary Report # 2335

$L_{\infty}$ -UPPER BOUND OF  $L_2$ -PROJECTIONS  
ONTO SPLINES AT A GEOMETRIC MESH

Rong-qing Jia

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

February 1982

(Received November 5, 1981)

DTIC FILE COPY

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

Approved for public release  
Distribution unlimited

DTIC  
ELECTE  
MAY 18 1982  
S D E

82 05 18 036

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

$L_\infty$ -UPPER BOUND OF  $L_2$ -PROJECTIONS  
ONTO SPLINES AT A GEOMETRIC MESH

Rong-qing Jia

Technical Summary Report #2335

February 1982

ABSTRACT

For an integer  $k > 1$  and a geometric mesh  $(q^i)_{-\infty}^{\infty}$  with  $q \in (0, \infty)$ , let

$$M_{i,k}(x) := k[q^i, \dots, q^{i+k}](-x)_+^{k-1}$$

$$N_{i,k}(x) := (q^{i+k} - q^i)M_{i,k}(x)/k,$$

and let  $A_k(q)$  be the Gram matrix  $(\int M_{i,k} N_{j,k})_{i,j \in \mathbb{Z}}$ . It is known that  $\|A_k(q)^{-1}\|_\infty$  is bounded independent of  $q$ . In this paper it is shown that  $\|A_k(q)^{-1}\|_\infty$  is strictly decreasing for  $q$  in  $[1, \infty)$ . In particular, the sharp upper bound and lower bound for  $A_k(q)^{-1}$  are obtained:

$$2k-1 < \|A_k(q)^{-1}\|_\infty < \left(\frac{\pi}{2}\right)^{2k} \left\{ \sum_{j \in \mathbb{Z}} (1+2j)^{-2k} \right\}^{-1} \text{ for all } q \in (0, \infty).$$

AMS (MOS) Subject Classification: 41A15

Key Words: splines, geometric mesh, least-squares approximation,

Gram matrix, monotonicity, sharp upper bound.

Work Unit Number 3 - Numerical Analysis and Computer Science

# SIGNIFICANCE AND EXPLANATION

Least-squares approximation by polynomial splines is a very effective means of approximation, particularly when the knot sequence can be chosen suitably nonuniform. The stability of this process can be linked to the norm of the inverse of the Gram matrix of a B-spline basis. For the special case when the knot sequence is geometric it is known that the norm of the inverse of that Gramian is bounded independently of the mesh ratio. Also, the sharp lower bound for the inverse of that Gramian is known.

In the present report, we continue these investigations and obtain, in particular, the sharp upper bound of the inverse of the Gram matrix.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

$L_\infty$ -UPPER BOUND OF  $L_2$ -PROJECTIONS ONTO SPLINES  
AT A GEOMETRIC MESH

Rong-qing Jia

1. Introduction

Let  $\underline{x} := (x_i)_{i=-\infty}^{\infty}$  be a strictly increasing biinfinite sequence with  $x_{\pm\infty} := \lim_{i \rightarrow \pm\infty} x_i$  and  $I := (x_{-\infty}, x_{+\infty})$ . Let further

$$S := \mathcal{M}_{k,\underline{x}}(I) := \{f \in C^{k-2}(I) \cap L_\infty(I) : f|_{(x_k, x_{i+1})} \text{ is a polynomial of degree } < k\}$$

be the normed linear space of bounded polynomial splines of order  $k$  with breakpoint sequence  $\underline{x}$  and norm  $\|f\| := \sup_{x \in I} |f(x)|$ . We shall be concerned with  $P_S$ , the orthogonal projector onto  $S$  with respect to the ordinary inner product

$$(f, g) := \int_I f(x)g(x)dx,$$

but restricted to  $L_\infty(I)$ . We want to bound its norm

$$\|P_S\|_\infty := \sup_{f \in L_\infty(I)} \|P_S f\|_\infty / \|f\|_\infty.$$

In 1973, de Boor raised the following

Conjecture [1].  $\sup_{\underline{x}} \|P_S\|_\infty < \text{const}_k < \infty.$

This conjecture has been verified for  $k = 1, 2, 3, 4$  (see de Boor [3] and the references cited there). de Boor [2] also obtained a bound of  $P_S$  in terms of a global mesh ratio. In general, however, this conjecture seems hard to solve. For geometric mesh  $\underline{x}$ , Höllig, K. [8] recently proved the boundedness of  $P_S$ . Later on, Feng, Y. Y. and Kozak, J. [6] reproved this result. Before recalling some results of theirs, we need to introduce some notations. For the mesh  $\underline{x} = (x_i)_{i=-\infty}^{\infty}$ , let

$$\begin{aligned} M_{i,k}(x) &:= k[x_1, \dots, x_{i+k}] (\cdot - x)_+^{k-1} \\ N_{i,k}(x) &:= ([x_{i+1}, \dots, x_{i+k}] - [x_1, \dots, x_{i+k-1}]) (\cdot - x)_+^{k-1} \\ &= (x_{i+k} - x_i) M_{i,k}(x) / k. \end{aligned}$$

Set

$$A_k(i, j) := \int M_{i, k} N_{j, k} \quad \text{for } i, j \in \mathbb{Z}.$$

Let  $A_k \in \mathbb{R}^{\mathbb{Z} \times \mathbb{Z}}$  be the biinfinite matrix given by the rule:

$$(i, j) \rightarrow A_k(i, j) \quad \text{for } (i, j) \in \mathbb{Z} \times \mathbb{Z}.$$

It was shown by de Boor [1] that

$$D_k^{-2} \|A_k^{-1}\|_{\infty} < \|P_S\|_{\infty} < \|A_k^{-1}\|_{\infty},$$

where  $D_k$  is a constant depending only on  $k$ . Thus bounding  $P_S$  is equivalent to bounding  $A_k^{-1}$ .

Let us restrict ourselves now to a particular case where  $\underline{x}$  is a geometric mesh:  
 $\underline{x} := (q^i)_{i=-\infty}^{\infty}$  for some  $q \in [1, \infty)$  (note that the case  $q \in (0, 1]$  is symmetric to the case  $q \in [1, \infty)$ ; see [6] and [8]). Spline interpolation at a geometric mesh was first investigated by C. A. Micchelli [9], who based his argument on the properties of the so-called generalized Euler-Frobenius polynomials. Later on, Y. Y. Feng and J. Kozak [6] developed such a consideration. Earlier, and in a different way, K. Höllig [8] made a more precise investigation into the boundedness of  $L_2$ -projections onto splines on a geometric mesh. In particular, he got the following elegant result (see Theorem 5 in [8]):

Theorem A. For a geometric mesh  $\underline{x} := (q^i)_{i=-\infty}^{\infty}$  with  $q \in (0, \infty)$ , let  $A_k(q)$  be the biinfinite matrix  $(\int M_{i, k} N_{j, k})_{i, j \in \mathbb{Z}}$ . Then

$$(1) \quad \|A_k(q)^{-1}\|_{\infty} = |\Omega_k(q)|^{-1},$$

where

$$(2) \quad \Omega_k(q) := 2 \cdot k! \cdot (k-1)! \cdot t^{2k-1} \prod_{v=1}^k \frac{q^{v+1}}{q^v - 1} \cdot \prod_{v=1}^{k-1} \frac{q^{v+1}}{q^v - 1} \cdot \sum_{j \in \mathbb{Z}} \prod_{v=1}^k \frac{1}{[\pi(1+2j)]^2 + (vt)^2}$$

with  $t := \log q$ . Moreover,

$$(3) \quad \lim_{q \rightarrow 1} \Omega_k(q) = \left(\frac{2}{\pi}\right)^{2k} \left\{ \sum_{j \in \mathbb{Z}} (1+2j)^{-2k} \right\}$$

$$(4) \quad \lim_{q \rightarrow \infty} \Omega_k(q) = \frac{1}{2k-1}.$$

Based on numerical evidence, de Boor raised the following

Conjecture.  $\Omega_k(q)$  is a monotone increasing function on  $[1, \infty)$ .

This conjecture was verified for  $k \leq q$  by Y. Y. Feng and J. Kozak [7]. They also showed that  $\Omega_k(q) \leq \frac{1}{2k-1}$  in the same paper.

The purpose of this paper is to confirm the above conjecture. Thus we have

Theorem 1.  $\Omega_k(q)$  is a monotone increasing function on  $[1, \infty)$ . In particular,

$$(5) \quad 2k-1 < \|A_k(q)^{-1}\|_{\infty} < \left(\frac{\pi}{2}\right)^{2k} \left\{ \sum_{j \in \mathbb{Z}} (1+2j)^{-2k} \right\}^{-1}.$$

Note that  $\Omega_1(q) \equiv 1$  and that  $\Omega_2(q) \equiv \frac{1}{3}$  in terms of a straightforward calculation. Hence we can restrict ourselves to the case  $k \geq 3$  from now on.

In section 2, we will give an alternative proof of theorem A. Section 3 and 4 will be devoted to proving the monotonicity of  $\Omega_k(q)$  for  $q \in [1, 20]$  and  $q \in [20, \infty)$ , respectively.

## 2. The bound for $A_k(q)^{-1}$ .

As before,  $\underline{x} = (q^i)_{i=0}^{\infty}$  is a geometric mesh with  $q \in (1, \infty)$  and  $t = \log q$ . Consider

$$\phi_0(x) := [0, 1, \dots, (2k-1)]_z \frac{x^z}{q^z + q^k} \text{ for } x \in [1, q].$$

It is easy to verify that

$$(6) \quad q^l \phi_0^{(l)}(q) + q^k \phi_0^{(l)}(1) = [0, 1, \dots, (2k-1)]_z \{z(z-1) \cdots (z-l+1)\} = \begin{cases} 0 & \text{for } l = 1, \dots, 2k-2 \\ 1 & \text{for } l = 2k-1 \end{cases}.$$

Since  $\phi_0$  is a polynomial of degree  $2k-1$ ,  $\phi_0^{(2k-1)}$  is constant in  $[1, q]$ . Hence (6) yields that

$$(7) \quad \phi_0^{(2k-1)}(x) = \frac{1}{q^k + q^{2k-1}} \text{ for } x \in [1, q].$$

Now we extend the domain of  $\phi_0$  to  $(0, \infty)$  as follows:

$$\phi(x) := (-q^k)^m \phi_0(q^{-m}x) \text{ for } q^m < x < q^{m+1}, m \in \mathbb{Z}.$$

From (6) we assert that  $\phi \in \mathcal{F}_{2k, \underline{x}}$ , and that

$$(8) \quad \phi(q^m) = (-q^k)^m \phi_0(1), m \in \mathbb{Z}.$$



It follows that

$$\begin{aligned} [x_0, x_1, \dots, x_{m-1}, x_m] \phi &= \frac{[x_1, \dots, x_m] \phi - [x_0, \dots, x_{m-1}] \phi}{x_m - x_0} \\ &= \frac{-q^{k-m+1} [x_0, \dots, x_{m-1}] \phi - [x_0, \dots, x_{m-1}] \phi}{q^m - 1} \\ &= -\frac{q^{k-m+1} + 1}{q^m - 1} [x_0, \dots, x_{m-1}] \phi. \end{aligned}$$

By induction on  $m$ , we can obtain

$$(9) \quad [x_0, x_1, \dots, x_k] \phi = (-1)^k \left( \prod_{m=1}^k \frac{q^{k-m+1} + 1}{q^m - 1} \right) \phi_0(1) = (-1)^k \left( \prod_{m=1}^k \frac{q^m + 1}{q^m - 1} \right) \cdot \phi_0(1).$$

From (8) we deduce that

$$(10) \quad [x_1, \dots, x_{i+k}] \phi = (-1)^i [x_0, \dots, x_k] \phi.$$

By Peano's theorem (see [4])

$$[x_1, \dots, x_{i+k}] \phi = \int M_{i,k}(x) \phi^{(k)}(x) / k! \, dx.$$

Now we get

$$(11) \quad \int M_{i,k}(x) \phi^{(k)}(x) / k! \, dx = (-1)^i (-1)^k \left( \prod_{m=1}^k \frac{q^m + 1}{q^m - 1} \right) \phi_0(1).$$

Obviously,  $\phi^{(k)} / k! \in \mathcal{S}_{k, \underline{x}}$ ; hence  $\phi^{(k)} / k!$  may be expanded in a B-spline series:

$$\phi^{(k)} / k! = \sum_j \alpha_j N_{j,k}.$$

However,  $\phi^{(k)}(qx) = -\phi^{(k)}(x)$ . Thus

$$\sum_j \alpha_j N_{j,k}(x) = -\sum_j \alpha_j N_{j,k}(qx) = -\sum_j \alpha_j N_{j-1,k}(x) = -\sum_j \alpha_{j+1} N_{j,k}(x).$$

By the uniqueness of B-spline expansion we assert that

$$\alpha_{j+1} = -\alpha_j, \quad j \in \mathbb{Z}.$$

Thus we can write

$$(12) \quad \phi^{(k)} / k! = C \sum_j (-1)^j N_{j,k},$$

where  $C$  is a constant to be determined. Now (11) and (12) together give

$$(13) \quad \sum_{j \in \mathbb{Z}} (-1)^j \int M_{i,k}(x) N_{j,k}(x) dx = (-1)^i C^{-1} (-1)^k \left( \prod_{m=1}^k \frac{q^m + 1}{q^m - 1} \right) \phi_0(1).$$

Let

$$(14) \quad \Omega_k(q) := C^{-1} (-1)^k \left( \prod_{m=1}^k \frac{q^m + 1}{q^m - 1} \right) \phi_0(1).$$

Then (see de Boor, C., S. Friedland and A. Pinkus [5])

$$\|A_k(q)^{-1}\|_\infty = |Q_k(q)|^{-1}.$$

It remains to determine C. Differentiate (12)  $k-1$  times:

$$\phi^{(2k-1)}/k! = C(\Sigma(-1)^j N_{j,k})^{(k-1)}.$$

On the one hand,

$$\phi^{(2k-1)}(x)/k! = \frac{1}{k!} \cdot \frac{1}{q^k + q^{2k-1}} \text{ for } x \in (1, q).$$

On the other hand (see [4]),

$$\begin{aligned} (\Sigma (-1)^j N_{j,k})^{(k-1)} &= \frac{2(k-1)}{q^{k-1}-1} \cdot \frac{(q+1)(k-2)}{q^{k-2}-1} \cdots \frac{q^{k-2}+1}{q-1} \cdot \sum_{j \in \mathbb{Z}} (-1)^j (q^{k-1})^{-j} N_{j,1} \\ &= 2 \cdot (k-1)! \cdot \frac{1}{q^{k-1}+1} \cdot \prod_{m=1}^{k-1} \frac{q^m+1}{q^m-1} \cdot \sum_{j \in \mathbb{Z}} (-1)^j (q^{k-1})^{-j} N_{j,1}. \end{aligned}$$

Thus, for  $x \in (1, q)$

$$(\Sigma (-1)^j N_{j,k})^{(k-1)}(x) = 2 \cdot (k-1)! \cdot \frac{1}{q^{k-1}+1} \cdot \prod_{m=1}^{k-1} \frac{q^m+1}{q^m-1}.$$

From the above calculation we get

$$\begin{aligned} (15) \quad C^{-1} &= k! \cdot (q^k + q^{2k-1}) \cdot 2 \cdot (k-1)! \cdot \frac{1}{q^{k-1}+1} \cdot \prod_{m=1}^{k-1} \frac{q^m+1}{q^m-1} \\ &= 2 \cdot k! \cdot (k-1)! \cdot q^k \prod_{m=1}^{k-1} \frac{q^m+1}{q^m-1}. \end{aligned}$$

Finally, (14) and (15) yield that

$$\begin{aligned} (16) \quad Q_k(q) &= (-1)^k \cdot 2 \cdot k! \cdot (k-1)! \prod_{m=1}^k \frac{q^m+1}{q^m-1} \cdot \prod_{m=1}^{k-1} \frac{q^m+1}{q^m-1} \cdot q^k \phi_0(1) \\ &= (-1)^k \cdot 2 \cdot k! \cdot (k-1)! \prod_{m=1}^k \frac{q^m+1}{q^m-1} \cdot \prod_{m=1}^{k-1} \frac{q^m+1}{q^m-1} \cdot q^k [0, 1, \dots, 2k-1] \frac{1}{q^{2k-1}+q^k}. \end{aligned}$$

We follow the procedure in [9] and use a well-known formula for the divided difference to get

$$(17) \quad (-1)^k q^k [0, 1, \dots, 2k-1] \frac{1}{q^{2k-1}+q^k} = \frac{(-1)^k q^k}{2\pi i} \left[ \int_{C_R} - \sum_{j=0}^{2k-1} \int_{C_j} \frac{dz}{\prod_{m=0}^{2k-1} (z-m)} \right] \cdot (e^{z \log q + q^k})$$

where  $C_R$  and  $C_{r_j}$  stand for positively oriented circles with centers at 0 and  $j$  and radius  $R$  and  $r_j$  where  $R$  sufficiently large and  $r_j$  sufficiently small,  $j = 0, 1, \dots, 2k-1$ .

Making  $R \rightarrow \infty$ ,  $r_j \rightarrow 0$  ( $j = 0, 1, \dots, 2k-1$ ) in (17) and using the residue theorem we get

$$\begin{aligned} (-1)^k q^k [0, 1, \dots, 2k-1] \frac{1}{q + q^k} &= (-1)^k q^k \cdot (-1) \sum_{j \in \mathbb{Z}} \operatorname{Res}_{z=i(\pi+2\pi j)/t+k} \frac{1}{\prod_{v=0}^{2k-1} (z-v) \cdot (e^{z \log q} + q^k)} \\ &= (-1)^k \cdot \sum_{j \in \mathbb{Z}} \frac{1}{t \prod_{v=0}^{2k-1} \left[ \frac{i(\pi+2\pi j)}{t} - v+k \right]} = \sum_{j \in \mathbb{Z}} \frac{t^{2k-1}}{\prod_{v=0}^{2k-1} [\pi+2\pi j - i(k-v)t]} \\ &= t^{2k-1} \cdot \sum_{j \in \mathbb{Z}} \prod_{v=1}^k \frac{1}{(\pi+2\pi j)^2 + (vt)^2}. \end{aligned}$$

Thus (2) is proved by substituting the above equality into (16). Then it is straightforward to verify (3). As to (4), we have

$$\begin{aligned} \lim_{q \rightarrow \infty} \Omega_k(q) &= (-1)^k \cdot 2 \cdot k! \cdot (k-1)! / (2k-1)! \cdot \lim_{q \rightarrow \infty} \sum_{l=0}^{2k-1} (-1)^{l+1} \binom{2k-1}{l} \frac{q^k}{q^l + q^k} \\ &= 2 \cdot (-1)^k / \binom{2k-1}{k} \cdot \left\{ \sum_{l=0}^{k-1} (-1)^{l+1} \binom{2k-1}{l} + (-1)^{k+1} \cdot \frac{1}{2} \cdot \binom{2k-1}{k} \right\} \\ &= 2 \cdot (-1)^k / \binom{2k-1}{k} \left\{ \sum_{l=0}^{k-1} (-1)^{l+1} \left[ \binom{2k-2}{l-1} + \binom{2k-2}{l} \right] + (-1)^{k+1} \cdot \frac{1}{2} \cdot \binom{2k-1}{k} \right\} \\ &= 2 \cdot (-1)^k / \binom{2k-1}{k} \left\{ (-1)^k \binom{2k-2}{k-1} + (-1)^{k+1} \cdot \frac{1}{2} \binom{2k-1}{k} \right\} = \frac{2k}{2k-1} - 1 = \frac{1}{2k-1}. \end{aligned}$$

This ends the proof of Theorem A.

### 3. The monotonicity of $\Omega_k(q)$ for $q \in [1, 20]$ .

Recall  $t = \log q$ . Let

$$f_k(t) := t^{2k-1} \cdot \prod_{v=1}^k \frac{e^{vt} + 1}{e^{vt} - 1} \cdot \prod_{v=1}^{k-1} \frac{e^{vt} + 1}{e^{vt} - 1} \cdot \sum_{j \in \mathbb{Z}} \prod_{v=1}^k \frac{1}{(\pi+2\pi j)^2 + (vt)^2}.$$

Then  $\Omega_k(e^t) = 2 \cdot k! (k-1)! f_k(t)$ . Consider  $f'_k(t)/f_k(t)$ . We have

$$(18) \quad \frac{f'_k(t)}{f_k(t)} = \sum_{j=0}^{\infty} \frac{1}{t} u_{k,j}(t) f_{k,j}(t) ,$$

where

$$u_{k,j}(t) := \prod_{v=1}^k \frac{1}{(\pi + 2\pi j)^2 + (vt)^2} / \left\{ \sum_{j=0}^{\infty} \prod_{v=1}^k \frac{1}{(\pi + 2\pi j)^2 + (vt)^2} \right\}$$

$$(19) \quad f_{k,j}(t) := 2k-1 + \sum_{v=1}^{k-1} + \sum_{v=1}^k \left( \frac{vte^{vt}}{e^{vt}+1} - \frac{vte^{vt}}{e^{vt}-1} \right) - \sum_{v=1}^k \frac{2(vt)^2}{(\pi + 2\pi j)^2 + (vt)^2} .$$

If we can show that  $f'_k(t)/f_k(t) > 0$  for  $t \in [0,3]$ , then  $\Omega'_k(q) > 0$  for  $q \in [1,20]$ , because  $e^3 > 20$ . For this it suffices to show  $f_{k,0}(t) > 0$ , since  $f_{k,j}(t) > f_{k,0}(t)$  ( $j = 1,2,\dots$ ) from (19). Let us first make the following observation.

Proposition 1.

$$\frac{\pi^2}{\pi^2 + (cx)^2} > \frac{2xe^x}{e^{2x}-1} \text{ for } x \in (0,\infty) \text{ and } c \in [1,5/4] .$$

Proof. Each of the following inequalities is equivalent to proposition 1:

$$e^{2x} - 1 > (1 + c^2 x^2 / \pi^2) 2xe^x ,$$

$$\sum_{n=0}^{\infty} \frac{2^n x^n}{(n+1)!} > (1 + c^2 x^2 / \pi^2) \left( \sum_{n=0}^{\infty} \frac{x^n}{n!} \right) ,$$

$$\sum_{n=2}^{\infty} \frac{2^n x^n}{(n+1)!} > \sum_{n=2}^{\infty} \left[ \frac{1}{n!} + \frac{c^2}{\pi^2} \cdot \frac{1}{(n-2)!} \right] x^n .$$

However, an induction argument on  $n$  shows that

$$\frac{2^n}{(n+1)!} > \frac{1}{n!} + \frac{c^2}{\pi^2} \cdot \frac{1}{(n-2)!} \text{ for } n \geq 2 \text{ and } c \in [1,5/4] .$$

Therefore proposition 1 is true.

Proposition 2.

$$\frac{2\pi^2}{\pi^2 + (4x/3)^2} > \frac{\pi^2}{\pi^2 + (5x/4)^2} + \frac{\pi^2}{\pi^2 + (5x/3)^2} .$$

Proof.

$$2(\pi^2 + 25x^2/9)(\pi^2 + 25x^2/16) = 2\pi^4 + 1250\pi^2 x^2/144 + 625x^4/72$$

$$> 2\pi^4 + 1137\pi^2 x^2/144 + 625x^4/81 = (\pi^2 + 16x^2/9)[(\pi^2 + 25x^2/9) + (\pi^2 + 25x^2/16)] .$$

Multiplying the above inequality by  $\pi^2 / [(\pi^2 + \frac{16}{9}x^2)(\pi^2 + \frac{25}{16}x^2)(\pi^2 + \frac{25}{9}x^2)]$ , we obtain proposition 2.

Proposition 3.  $f_{k+1,0}(t) > f_{k,0}(t)$  for  $t > 0$  and  $k > 3$ .

Proof. We shall argue by induction on  $k$ . For  $k = 3$ , we have

$$f_{4,0}(t) - f_{3,0}(t) = \frac{2\pi^2}{\pi^2 + (4t)^2} - \frac{2 \cdot 3t \cdot e^{3t}}{e^{2 \cdot 3t} - 1} - \frac{2 \cdot 4t \cdot e^{4t}}{e^{2 \cdot 4t} - 1}.$$

Set  $x := 3t$ . Then proposition 1 and 2 yield that

$$f_{4,0}(t) - f_{3,0}(t) > \left( \frac{\pi^2}{\pi^2 + (5x/4)^2} - \frac{2xe^x}{e^{2x} - 1} \right) + \left( \frac{\pi^2}{\pi^2 + (5x/3)^2} - \frac{2 \cdot \frac{4}{3}xe^{4x/3}}{e^{2 \cdot 4x/3} - 1} \right) > 0$$

Suppose now that  $k > 4$ . Then  $\frac{k+1}{k} < \frac{5}{4}$ . We have

$$\begin{aligned} f_{k+1,0}(t) - f_{k,0}(t) &= \frac{2\pi^2}{\pi^2 + [(k+1)t]^2} - \frac{2 \cdot kt \cdot e^{kt}}{e^{2kt} - 1} - \frac{2(k+1)t e^{(k+1)t}}{e^{2(k+1)t} - 1} \\ &= \left( \frac{\pi^2}{\pi^2 + [(k+1)t]^2} - \frac{2 \cdot kt \cdot e^{kt}}{e^{2kt} - 1} \right) + \left( \frac{\pi^2}{\pi^2 + [(k+1)t]^2} - \frac{2 \cdot (k+1)t \cdot e^{(k+1)t}}{e^{2 \cdot (k+1)t} - 1} \right) > 0, \end{aligned}$$

according to proposition 1. Thus proposition 3 is proved.

Consequently,  $f_{k,j}(t) > f_{3,0}(t)$  for all  $k > 3$  and  $j > 0$ . The remaining task of this section is to elaborate the nonnegativity of  $f_{3,0}(t)$ . For this we need some estimates.

Proposition 4. Let  $h(x) = \left( \frac{xe^x}{e^x + 1} - \frac{1}{2}x \right) \cdot \frac{1}{x}$ . Then  $h'(x) < 0$  for  $x > 0$ .

Proof.  $h(x) = \frac{e^x - 1}{2x(e^x + 1)}$  and

$$h'(x) = \frac{1}{2} \cdot \frac{x(e^x + 1)e^x - (e^x - 1)[(e^x + 1) + xe^x]}{[x(e^x + 1)]^2} = \frac{1 + 2xe^x - e^{2x}}{2x^2(e^x + 1)^2},$$

while

$$1 + 2xe^x - e^{2x} = 1 + 2x \cdot \sum_{n=0}^{\infty} \frac{x^n}{n!} - \sum_{n=0}^{\infty} \frac{2^n x^n}{n!} = - \sum_{n=3}^{\infty} \frac{2(2^{n-1} - n)}{n!} x^n < 0, \text{ for } x > 0.$$

Proposition 5.  $-\frac{xe^x}{e^x - 1} > -1 - \frac{1}{2}x - \frac{1}{12}x^2$

$$\begin{aligned} \text{Proof. } (1 + \frac{1}{2}x + \frac{1}{12}x^2)(e^x - 1) - xe^x &= \sum_{n=2}^{\infty} \frac{x^{n+1}}{n!} - \sum_{n=1}^{\infty} \frac{1}{2} \cdot \frac{x^{n+1}}{n!} + \sum_{n=1}^{\infty} \frac{1}{12} \cdot \frac{x^{n+2}}{n!} \\ &= \sum_{n=5}^{\infty} \frac{1}{n!} \cdot \frac{n^2 - 7n + 12}{12} x^n > 0 \text{ for } x > 0. \end{aligned}$$

Now we are in a position to prove that  $f_{3,0}(t) > 0$  for  $t \in [0, 0.3]$ . Write down

$$\begin{aligned} f_{3,0}(t) &= 2\left(1 + \frac{te^t}{e^t+1} - \frac{te^t}{e^t-1} - \frac{t^2}{t^2+\pi^2}\right) + 2\left(\frac{2te^{2t}}{e^{2t}+1} - \frac{2te^{2t}}{e^{2t}-1} - \frac{(2t)^2}{(2t)^2+\pi^2}\right) \\ &\quad + \left(1 + \frac{3te^{3t}}{e^{3t}+1} - \frac{3te^{3t}}{e^{3t}-1} - \frac{(3t)^2}{(3t)^2+\pi^2}\right) - \frac{9t^2}{\pi^2+9t^2}. \end{aligned}$$

It follows from proposition 4 that, for  $t \in [0, 0.3]$ ,

$$te^t/(e^t+1) - t/2 > h(0.3) \cdot t^2 > 0.248t^2$$

$$2te^{2t}/(e^{2t}+1) - 2t/2 > h(0.6) \cdot (2t)^2 > 0.242 \cdot (2t)^2$$

$$3te^{3t}/(e^{3t}+1) - 3t/2 > h(0.9) \cdot (3t)^2 > 0.234 \cdot (3t)^2.$$

In connection with proposition 5, we obtain

$$2\left(1 + \frac{te^t}{e^t+1} - \frac{te^t}{e^t-1} - \frac{t^2}{t^2+\pi^2}\right) > 2\left(0.248 - \frac{1}{12} - \frac{1}{\pi^2}\right)t^2 > 0.126t^2,$$

$$2\left(1 + \frac{2te^{2t}}{e^{2t}+1} - \frac{2te^{2t}}{e^{2t}-1} - \frac{(2t)^2}{(2t)^2+\pi^2}\right) > 2\left(0.242 - \frac{1}{12} - \frac{1}{\pi^2}\right)(2t)^2 > 0.458t^2,$$

$$1 + \frac{3te^{3t}}{e^{3t}+1} - \frac{3te^{3t}}{e^{3t}-1} - \frac{(3t)^2}{(3t)^2+\pi^2} > \left(0.234 - \frac{1}{12} - \frac{1}{\pi^2}\right)(3t)^2 > 0.444t^2,$$

$$-\frac{9t^2}{\pi^2+9t^2} > -\frac{9t^2}{\pi^2} > -0.912t^2.$$

As a conclusion,

$$f_{3,0}(t) > (0.126 + 0.458 + 0.444 - 0.912)t^2 = 0.116t^2.$$

This shows that

$$f'_k(t) > 0 \text{ for } t \in [0, 0.3].$$

The next case we are going to treat is that  $t \in [0.3, 3]$ . Let

$$v(t) := \frac{2t^2}{t^2 + \pi^2} + \frac{2t^2}{4t^2 + \pi^2} + \frac{t^2 - 9t^2}{\pi^2 + 9t^2}$$

$$w(t) := \frac{4te^t}{e^{2t}-1} + \frac{8te^{2t}}{e^{4t}-1} + \frac{6te^{3t}}{e^{6t}-1}$$

Then  $f_{3,0}(t) = v(t) - w(t)$ . It is easily seen that  $v'(t) < 0$  for  $t \in [0, \infty)$ . We claim that  $w'(t) < 0$  ( $0 < t < \infty$ ), too. This is guaranteed by the following proposition.

Proposition 6. Let  $g(x) := \frac{2xe^x}{e^{2x}-1}$ . Then  $g'(x) < 0$  for  $x > 0$ .

Proof.  $g'(x) = -\frac{2e^x}{(e^{2x}-1)^2} (1 + x + xe^{2x} - e^{2x})$ , while

$$1 + x + xe^{2x} - e^{2x} = (1+x) - (1-x) \sum_{n=0}^{\infty} \frac{2^n x^n}{n!} = \sum_{n=3}^{\infty} \frac{(n-2) \cdot 2^{n-1}}{n!} x^n > 0 \text{ for } x > 0.$$

Accordingly,

$$(20) \quad f_{3,0}(t) = v(t) - w(t) > v(b) - w(a) \text{ for } t \in [a, b] \text{ with } 0 < a < b.$$

To determine the positivity of  $f_{3,0}$  I wrote a Fortran program and found that

$$v\left(\frac{n+1}{100}\right) - w\left(\frac{n}{100}\right) > 0.001 \text{ for } n = 30, 31, \dots, 299.$$

Thus by (20) we assert that

$$f_{3,0}(t) > 0 \text{ for } t \in \left[\frac{n}{100}, \frac{n+1}{100}\right], n = 30, 31, \dots, 299.$$

Therefore

$$f_{3,0}(t) > 0 \text{ for } t \in \bigcup_{n=30}^{299} \left[\frac{n}{100}, \frac{n+1}{100}\right] = [0.3, 3].$$

So far we have shown that  $\Omega'_k(q) > 0$  for  $q \in [1, 20]$ .

#### 4. The monotonicity of $\Omega_k(q)$ for $q \in [20, \infty)$ .

Let

$$(21) \quad f(q) := (-1)^k \cdot (2k-1)! q^k [0, 1, \dots, 2k-1] \frac{1}{q + q^k}.$$

Then

$$\begin{aligned} f(q) &= (-1)^k q^k \sum_{l=0}^{2k-1} \binom{2k-1}{l} (-1)^{l+1} \frac{1}{q + q^k} = (-1)^k \sum_{l=0}^{2k-1} (-1)^{l+1} \binom{2k-1}{l} \frac{1}{1+q^{k-l}} \\ &= \sum_{l=0}^{k-1} (-1)^{k+l+1} \binom{2k-1}{l} \frac{q^{k-l}}{1+q^{k-l}} + (-1)^k \binom{2k-1}{k} \frac{1}{2} + \sum_{l=k+1}^{2k-1} (-1)^{k+l+1} \binom{2k-1}{l} \frac{1}{1+q^{l-k}} \end{aligned}$$

It follows that

$$\begin{aligned}
 (22) \quad f'(q) &= \sum_{\ell=0}^{k-1} (-1)^{k+\ell+1} \binom{2k-1}{\ell} \frac{(k-\ell)q^{k-\ell-1}}{(1+q)^{k-\ell+1}} + \sum_{\ell=k+1}^{2k-1} (-1)^{k+\ell} \binom{2k-1}{\ell} \frac{(\ell-k)q^{\ell-k-1}}{(1+q)^{\ell-k+1}} \\
 &= \sum_{\ell=1}^k (-1)^{\ell-1} \binom{2k-1}{k-\ell} \frac{\ell q^{\ell-1}}{(1+q)^{\ell+1}} - \sum_{\ell=1}^{k-1} (-1)^{\ell-1} \binom{2k-1}{k+\ell} \frac{\ell q^{\ell-1}}{(1+q)^{\ell+1}} \\
 &= \sum_{\ell=1}^{k-1} (-1)^{\ell-1} \left[ \binom{2k-1}{k-\ell} - \binom{2k-1}{k+\ell} \right] \frac{\ell q^{\ell-1}}{(1+q)^{\ell+1}} + (-1)^{k-1} \frac{k q^{k-1}}{(1+q)^k}.
 \end{aligned}$$

Now we need the following propositions.

Proposition 7.  $\left[ \binom{2k-1}{k-\ell} - \binom{2k-1}{k+\ell} \right] \frac{\ell q^{\ell-1}}{(1+q)^{\ell+1}}$  decreases as  $\ell$  increases and  $q > 6$ .

Proof.

$$(23) \quad \binom{2k-1}{k-\ell} - \binom{2k-1}{k+\ell} = \frac{(2k-1)!}{(k-\ell)!(k+\ell)!} \left( 1 - \frac{k-\ell}{k+\ell} \right) = \frac{(2k-1)!}{(k-\ell)!(k+\ell)!} \cdot 2\ell.$$

We want to show

$$(24) \quad \frac{(2k-1)!}{(k-\ell)!(k+\ell)!} \cdot 2\ell \cdot \frac{\ell q^{\ell-1}}{(1+q)^{\ell+1}} > \frac{(2k-1)!}{(k-\ell-1)!(k+\ell+1)!} \cdot 2(\ell+1) \cdot \frac{(\ell+1)q^{\ell}}{(1+q)^{\ell+1}} \text{ for } q > 6.$$

It is easily seen that (24) is equivalent to

$$\frac{1}{q} \left( \frac{q^{\ell+1}+1}{q^{\ell}+1} \right)^2 > \frac{k-\ell}{k+\ell+1} \left( 1 + \frac{1}{\ell} \right)^2.$$

However,

$$\frac{1}{q} \left( \frac{q^{\ell+1}+1}{q^{\ell}+1} \right)^2 = \frac{q^{2\ell+2}+2q^{\ell+1}+1}{q(q^{2\ell}+2q^{\ell}+1)} > q - 2q^{1-\ell} > q-2,$$

because

$$(q-2q^{1-\ell})q(q^{2\ell}+2q^{\ell}+1) = q^{2\ell+2} - q^2 - 2q^{2-\ell} < q^{2\ell+2} + 2q^{\ell+1} + 1.$$

Meanwhile

$$\frac{k-\ell}{k+\ell+1} \left( 1 + \frac{1}{\ell} \right)^2 < 4.$$

Therefore (24) holds for  $q > 6$ , and proposition 7 is proved.

Proposition 8. For  $k \geq 2$  and  $q > 6$ ,

$$(25) \quad f'(q) > \binom{2k-1}{k-1} \cdot \frac{2}{k+1} \cdot \frac{1}{(1+q)^2} - \binom{2k-1}{k-2} \cdot \frac{4}{k+2} \cdot \frac{2q}{(1+q)^2}.$$

In particular,  $f'(q) > 0$  and

$$(26) \quad f(q) \leq \lim_{q \rightarrow \infty} f(q) = \binom{2k-1}{k-1} \cdot \frac{1}{2(2k-1)}.$$



Proof. Suppose first  $k$  is even,  $k = 2m$ . Then (22) and (23) yield that

$$\begin{aligned} f'(q) &= \binom{2k-1}{k-1} \cdot \frac{2}{k+1} \cdot \frac{1}{(1+q)^2} - \binom{2k-1}{k-2} \cdot \frac{4}{k+2} \cdot \frac{2q}{(1+q)^2} \\ &+ \sum_{j=2}^{m-1} \left[ \frac{(2k-1)!}{(k-2j+1)!(k+2j-2)!} \cdot \frac{2(2j-1)}{k+2j-1} \cdot \frac{(2j-1)q^{2j-2}}{(1+q)^{2j-1}} \right. \\ &\quad \left. - \frac{(2k-1)!}{(k-2j)!(k+2j-1)!} \cdot \frac{2 \cdot 2j}{k+2j} \cdot \frac{2jq^{2j-1}}{(1+q)^{2j}} \right] \\ &+ [(2k-2) \cdot \frac{(k-1)q^{k-2}}{(1+q)^{k-1}} - \frac{kq^{k-1}}{(1+q)^k}] . \end{aligned}$$

By proposition 7 all the terms under the summation sign are positive. Moreover,

$$q^{k-2}/(1+q)^{k-1} > q^{k-1}/(1+q)^k \text{ for } q > 1 ,$$

and

$$(2k-2)(k-1) \cdot \frac{q^{k-2}}{(1+q)^{k-1}} - \frac{kq^{k-1}}{(1+q)^k} > [(2k-2)(k-1) - k] \frac{q^{k-1}}{(1+q)^k} > 0 \text{ for } k > 2 .$$

For odd  $k$ , the proof is similar. Thus (25) holds. Furthermore

$$f'(q) > \frac{2}{(1+q)^2} \frac{(2k-1)!}{(k-2)!(k+2)!} \frac{k+2}{k-1} - \frac{4(1+1/q)^2}{q(1+1/q^2)^2} > \frac{2}{(1+q)^2} \frac{(2k-1)!}{(k-2)!(k+2)!} [1-4 \cdot (1 + \frac{1}{6})^2] \geq 0$$

for  $q > 6$  ,

and

$$f(q) \leq \lim_{q \rightarrow \infty} f(q) = \lim_{q \rightarrow \infty} \left[ \binom{2k-1}{k-1} \cdot \frac{1}{2} \cdot \Omega_k(q) \right] = \binom{2k-1}{k-1} \cdot \frac{1}{2(2k-1)} .$$

Proposition 9. Let  $S(q) = 4 \sum_{v=1}^{k-1} \frac{vq^{v-1}}{q^{2v-1}} + 2 \frac{kq^{k-1}}{q^{2k-1}}$ . Then

$$S(q) < \frac{4q^2}{(q^2-1)(q-1)^2} \text{ for } q > 1 .$$

Proof. We have

$$S(q) = 4 \cdot \sum_{v=1}^{k-1} \frac{q^{2v}}{q^{2v-1}} vq^{-(v+1)} + 2k \cdot q^{-(k+1)} \cdot \frac{q^{2k}}{q^{2k-1}} .$$

Note that

$$\frac{q^{2v}}{q^{2v-1}} < \frac{q^2}{q^2-1} \text{ for } v > 1 \text{ and } q > 1 .$$

Hence

$$S(q) \leq \frac{4q^2}{q^2-1} \cdot \sum_{v=1}^{\infty} vq^{-(v+1)} = \frac{4q^2}{q^2-1} \cdot \frac{d}{dq} \left( - \sum_{v=1}^{\infty} q^{-v} \right) = \frac{4q^2}{q^2-1} \cdot \frac{d}{dq} \left( \frac{-1}{1-q} \right) = \frac{4q^2}{(q^2-1)(q-1)^2}.$$

Proposition 10. Let  $g_k(q) = \frac{2k-1}{k+1} \left[ 1 - \frac{4(k-1)}{k+2} \cdot \frac{q}{1+q^2} \right]$ . Then

$$g_{k+1}(q) > g_k(q) \quad (k = 1, 2, 3, \dots; q > 12).$$

Proof.

$$g_{k+1}(q) - g_k(q) = \frac{3}{(k+1)(k+2)(k+3)} [(k+3) - (3k-1) \cdot 4 \cdot \frac{q}{1+q^2}]$$

$$> \frac{3}{(k+1)(k+2)(k+3)} [(k+3) - (3k-1) \cdot 4 \cdot \frac{12}{1+12^2}] > 0 \quad \text{for } q > 12.$$

Now we are in a position to prove the monotonicity of  $\Omega_k(q)$  for  $q \in [20, \infty)$ . From (16) and (21) we see that

$$\Omega_k(q) = 2^k k! (k-1)! / (2k-1)! \cdot \prod_{m=1}^k \frac{q^{m+1}}{q^m-1} \cdot \prod_{m=1}^{k-1} \frac{q^{m+1}}{q^m-1} \cdot f(q).$$

Hence

$$\frac{\Omega'_k(q)}{\Omega_k(q)} = - \left( 4 \sum_{v=1}^{k-1} \frac{vq^{v-1}}{q^{2v}-1} + 2 \frac{kq^{k-1}}{q^{2k}-1} \right) + \frac{f'(q)}{f(q)} = -S(q) + \frac{f'(q)}{f(q)}.$$

By proposition 9 we have

$$S(q) \leq \frac{1}{q^2} \cdot \frac{4q^4}{(q^2-1)(q-1)^2} \leq \frac{1}{q^2} \cdot \frac{4 \cdot 20^4}{(20^2-1)(20-1)^2} \leq \frac{4.45}{q^2} \quad \text{for } q > 20.$$

Moreover, proposition 8 and 10 tell us that

$$\begin{aligned} \frac{f'(q)}{f(q)} &> \frac{1}{(1+q)^2} \cdot \frac{4(2k-1)}{k+1} \cdot \left[ 1 - \frac{4(k-1)}{k+2} \cdot \frac{q}{1+q^2} \right] = \frac{4}{(1+q)^2} g_k(q) \\ &> \frac{4}{(1+q)^2} g_4(q) = \frac{4}{(1+q)^2} \cdot \frac{7}{5} \cdot \left( 1 - \frac{2q}{1+q^2} \right) = \frac{1}{q^2} \cdot \frac{28}{5} \frac{(1-1/q)^2}{(1+1/q)^2} \cdot \frac{1}{1+1/q^2} \\ &> \frac{1}{q^2} \cdot \frac{28}{5} \cdot \frac{(1-\frac{1}{20})^2}{(1+\frac{1}{20})^2} \cdot \frac{1}{1+1/20^2} > \frac{4.55}{q^2} \quad \text{for } q > 20 \text{ and } k > 4. \end{aligned}$$

Therefore

$$\frac{\Omega'_k(q)}{\Omega_k(q)} = \frac{f'(q)}{f(q)} - S(q) > \frac{4.55}{q^2} - \frac{4.45}{q^2} = \frac{0.1}{q^2} > 0 \quad \text{for } k > 4 \text{ and } q > 20.$$

It remains to check the case  $k = 3$ . For this we shall make a straightforward computation:

$$\begin{aligned}
\Omega_3(q) &= 24 \cdot \left(\frac{q+1}{q-1}\right)^2 \cdot \left(\frac{q^2+1}{q^2-1}\right)^2 \cdot \frac{q^3+1}{q^3-1} \cdot q^3 \cdot (-1)^{[0,1,2,3,4,5]} \cdot \frac{1}{q^2+q^3} \\
&= \frac{24}{120} \left(\frac{q+1}{q-1}\right)^2 \cdot \left(\frac{q^2+1}{q^2-1}\right)^2 \cdot \frac{q^3+1}{q^3-1} \cdot q^3 \left( \frac{1}{1+q^3} - 5 \frac{1}{q+q^3} + 10 \frac{1}{q^2+q^3} - 10 \cdot \frac{1}{2q^3} + 5 \frac{1}{q^3+q^4} - \frac{1}{q^3+q^5} \right) \\
&= \frac{1}{5} \cdot \frac{q^2+1}{q^2+q+1} .
\end{aligned}$$

Thus

$$\Omega'_3(q) = \frac{1}{5} \cdot \frac{q^2-1}{(q^2+q+1)^2} > 0 \text{ for } q > 1 .$$

This completes the proof of theorem 1.

#### Acknowledgement

I am glad to thank Professor Carl de Boer for his valuable help. Especially, section 2 and 4 reflect on his suggestion.

# REFERENCES

- [1] de Boor, C., The quasi-interpolant as a tool in elementary polynomial spline theory, in "Approximation Theory," G. G. Lorentz, ed., Academic Press (1973), 269-276.
- [2] de Boor, C., A bound on the  $L_\infty$ -norm of  $L_2$ -approximation by splines in terms of a global mesh ratio, Math. Comp. 30 (1976), 765-771.
- [3] de Boor, C., On a max-norm bound for the least squares splines approximant, in "Approximation and Function Spaces," Z. Ciesielski, ed.
- [4] de Boor, C., A practical guide to splines, Springer-Verlag, New York, 1978.
- [5] de Boor, C., S. Friedland and A. Pinkus, Inverses of infinite sign regular matrices, MRC TSK 2159 (1980); Trans. Amer. Math. Soc., to appear.
- [6] Peng, Y. Y. and J. Kozak, On the generalized Euler-Frobenius polynomial, MRC TSR #2088, J. Approximation Theory, to appear.
- [7] Peng, Y. Y. and J. Kozak,  $L_\infty$ -lower bound of  $L_2$ -projections onto splines on a geometric mesh, MRC TSR #2178, J. Approximation Theory, to appear.
- [8] Höllig, K.,  $L_\infty$ -boundedness of  $L_2$ -projections on splines for a geometric mesh, J. Approximation Theory, to appear.
- [9] Micchelli, C. A., Cardinal L-splines, in "Studies in Spline Functions and Approximation Theory," Academic Press (1976), 203-250.

RqJ/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2335	2. GOVT ACCESSION NO. AD-A274 697	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  $L_\infty$ -Upper Bound of $L_2$ -Projections onto Splines at a Geometric Mesh		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  Rong-qing Jia		8. CONTRACT OR GRANT NUMBER(s)  DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Computer Science
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE February 1982
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 15
		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION, DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  splines, geometric mesh, least-squares approximation, Gram matrix, monotonicity, sharp upper bound.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) For an integer $k \geq 1$ and a geometric mesh $(q^i)_{-\infty}^{\infty}$ with $q \in (0, \infty)$ , let $M_{i,k}(x) := k[q^i, \dots, q^{i+k}](-x)^{k-1}_+$ $N_{i,k}(x) := (q^{i+k} - q^i)M_{i,k}(x)/k,$ and let $A_k(q)$ be the Gram matrix $(\int M_{i,k} N_{j,k})_{i,j \in \mathbb{Z}}$ . It is known that <div style="text-align: right;">(continued)</div>		

ABSTRACT (continued)

$\|A_k(q)^{-1}\|_\infty$  is bounded independent of  $q$ . In this paper it is shown that  $\|A_k(q)^{-1}\|_\infty$  is strictly decreasing for  $q$  in  $[1, \infty)$ . In particular, the sharp upper bound and lower bound for  $A_k(q)^{-1}$  are obtained:

$$2k-1 \leq \|A_k(q)^{-1}\|_\infty \leq \left(\frac{\pi}{2}\right)^{2k} \left\{ \sum_{j \in \mathbb{Z}} (1+2^j)^{-2k} \right\}^{-1} \text{ for all } q \in (0, \infty) .$$

DATE  
ILME  
—8